

An Exact Expression of the Collective Excitation Energy Gap of Fractional Quantum Hall Effect

Yu-Liang Liu*

International Center for Theoretical Physics, P. O. Box 586, 34100 Trieste, Italy

Abstract

We have exactly solved the eigenequation of a two-dimensional Dirac fermion moving on the surface of a sphere under the influence of a radial magnetic field B , and obtained an exact expression of the collective excitation energy gap for the filling factors $\nu = \frac{p}{2mp \pm 1}$, m and p are non-zero integers, which is very well agreement with the computing results.

Recently, there have been much progress in understanding of the fractional quantum Hall effect(FQHE) [1], which results from a strongly correlated incompressible fluid state [2] formed at special densities n_e of a two-dimensional(2D) electron gas subject to a perpendicular magnetic field B . Based on the idea of "composite fermions", so called by Jain [3], resulting from the highly correlated motion of the carries, which effectively attaches even flux quanta to each electron[[4]- [8]], Halperin, Lee and Read [9] proposed a theory of a compressible Fermi-liquid-like state at the half-filled Landau level, $\nu = \frac{1}{2}$, which can be regarded as a degenerate system of the composite fermions in the absence of a magnetic field, populating a Fermi sea. If the filling factor differs from $\frac{1}{2}$, the composite fermions see a net magnetic field $\Delta B = B - B_{1/2}$, $B_{1/2} = 2n_e\Phi_0$ ($\Phi_0 = hc/e$ is the flux quantum), and at filling factor $\nu = p/(2p + 1)$, and p being a nonzero integer, they may fill $|p|$ Landau levels, and correspond to the incompressible quantized Hall states whose energy gaps are proportional to the net magnetic fields $\Delta B = n_e\Phi_0/p$. these predictions are strongly supported by the current experimental data[[10]- [19]].

On the other hand, there have been a lot of numerical calculations[[20]- [25]]employing the spherical geometry [26], in which N electrons move on the surface of a sphere under the influence of a radial magnetic field, to study the elementary excitations in fractional quantum Hall system. The flux through the surface of the sphere is $2S\Phi_0$, S is an integer or half-integer, the single electron states on the sphere have eigenfunctions $Y_{S,l,m}$, called "monopole harmonics", and eigenvalues[[27], [28]], $E_{l,m} = \hbar \frac{2l(l+1)-S^2}{2m_e R^2}$, where m_e is the electron band mass and R is the radius of the sphere, and satisfies the relation $4\pi R^2 B = 2S\Phi_0$. The qualitative agreement with the numerical results of interaction electrons at the magnetic field B and noninteracting fermions at the magnetic field ΔB gives a remarkable confirmation of the composite fermion picture. However, in all these approaches, there involves in the electron band mass m_e in their basic single quasiparticle excitations, but it is widely accepted that the FQHE derives from the strongly correlated interaction of the electrons within single Landau level, the basic energy unit of the quasiparticle excitation is e^2/l_B , $l_B = \sqrt{\hbar/(eB)}$ being the magnetic length. It is very difficult to scale the electron band mass m_e by this

basic length l_B [9]. In Ref. [29], we have given a simple explanation for the FQHE at even and odd denominator filling factors from the viewpoint of the topologic property of a 2D Dirac fermion gas, and used this topologic property to formally define a correspondence between the representations of the quantum Hall system in terms of electrons and composite fermions in the usual gauge field approaches[[4]- [8]]. The basic topologic term of the 2D Dirac fermion gas will automatically enforce a Dirac fermion to attach with two flux quanta, which becomes a basic composite fermion. In this Letter, we give an exact solution of a 2D Dirac fermion moving on the surface of a sphere under the influence of a radial magnetic field produced by a monopole at the centre of the sphere, and study the elementary excitation spectrum in the fractional quantum Hall system. Of course, the fractional quantum Hall effect derives from the strong Coulomb interaction of the conduction electrons, to treat it, we must consider this strong Coulomb interaction of the conduction electrons, however, up to now we have not found a reasonable method directly to treat it but the Laughlin's trial wavefunction. We think that the 2D Dirac fermion only describes a collective excitation mode of this strong Coulomb interaction of the conduction electrons within single Landau level, therefore there only exists a Coulomb-like energy scale e^2/l_B in the 2D Dirac fermion system, it is a basic energy unit which has not any direct relation with a real electron mass and the Coulomb interaction of the electrons. For a real material, its character is reflected only by the dielectric constant ϵ which enters the 2D Dirac fermion system with electric charge e . By using the 2D Dirac fermion to study the fractional quantum Hall effect, we can avoid directly to treat the strongly coupling electronic system.

For a 2D Dirac fermion system, its Hamiltonian can be written in the surface of a sphere as [31]

$$H = \int dx \bar{\psi} [-i\gamma^\mu D_\mu \pm m] \psi \quad (1)$$

where γ_μ are the 2D Dirac matrices, $\{\gamma_\mu, \gamma_\nu\} = \delta_{\mu\nu}$, $\mu, \nu = 1, 2$; $D_\mu = \partial_\mu - i(\Gamma_\mu + A_\mu)$, Γ_μ are the spin-connection, A_μ are the gauge field produced by the monopole at the centre of the sphere which is defined as $4\pi R^2 B = 2S\Phi_0$, R is the radius of the sphere, B is an

external magnetic field, $2S$ (an integer) is the strength of the monopole, m is the "mass" (chemical potential) of the 2D Dirac fermion which induces a topologic term (i.e., Chern-Simons term): $\pm \frac{1}{8\pi} \int d\tau d^2x \epsilon_{\mu\nu\lambda} a_\mu \partial_\nu a_\lambda$ in a plane space, a_μ are the Chern-Simons gauge fields [30], this Chern-Simons term enforces the 2D Dirac fermion attaching with two flux quanta [29], therefore the 2D Dirac fermion can be seen as an effective composite fermion. Here we will omit the mass term in (1) because it does not affect our final results. Due to in the 2D Dirac fermion gas, there naturally exists a topologic term which enforces the fermion to carry two flux quanta each [29], we can use this 2D Dirac fermion gas to mimic the strongly correlated electronic system of the FQHE, the 2D Dirac fermion should be considered in mind as a collective excitation mode produced by the strong Coulomb interaction of the conduction electrons within single Landau level, for simplicity, we omit the interaction among the 2D Dirac fermions. To avoid the singularities, we divide the surface of the sphere into two regions R_a and R_b , and define two gauge fields in these two regions, respectively [27]

$$\begin{aligned} A_r = A_\theta = 0, \quad A_\phi &= \frac{2S}{R \sin \theta} (1 - \cos \theta), \quad 0 \leq \theta < \frac{\pi}{2} + \delta, \quad 0 \leq \phi < 2\pi \\ A_r = A_\theta = 0, \quad A_\phi &= \frac{-2S}{R \sin \theta} (1 + \cos \theta), \quad \frac{\pi}{2} - \delta < \theta \leq \pi, \quad 0 \leq \phi < 2\pi \end{aligned} \quad (2)$$

where δ is a constant, $0 < \delta < \frac{\pi}{2}$. From Eq.(1), we can obtain the following eigenequation of single 2D Dirac fermion

$$-i \begin{pmatrix} 0, & \frac{A}{R} \\ \frac{\bar{A}}{R}, & 0 \end{pmatrix} \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} = E \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} \quad (3)$$

where $A = -i(\partial_\theta - \frac{i}{\sin \theta} \partial_\phi + \frac{\cos \theta}{2 \sin \theta} - RA_\phi)$, $\bar{A} = -i(\partial_\theta + \frac{i}{\sin \theta} \partial_\phi + \frac{\cos \theta}{2 \sin \theta} + RA_\phi)$. According to Eq.(3), we have the following equations

$$\begin{aligned} A\bar{A}\psi_1 &= R^2 E^2 \psi_1 \\ \bar{A}A\psi_2 &= R^2 E^2 \psi_2 \end{aligned} \quad (4)$$

Now we define two sets of angular momentum operators

$$\begin{cases} J^z = -i\partial_\phi - S' \\ J^\pm = \pm e^{\pm i\phi} \partial_\theta + i e^{\pm i\phi} \frac{\cos \theta}{\sin \theta} \partial_\phi + e^{\pm i\phi} \left(-\frac{2S-1}{2 \sin \theta} + \frac{S' \cos \theta}{\sin \theta} \right) \end{cases} \quad (5)$$

$$\begin{cases} L^z = -i\partial_\phi - S' \\ L^\pm = \pm e^{\pm i\phi} \partial_\theta + ie^{\pm i\phi} \frac{\cos \theta}{\sin \theta} \partial_\phi + e^{\pm i\phi} \left(-\frac{2S+1}{2\sin \theta} + \frac{S' \cos \theta}{\sin \theta} \right) \end{cases} \quad (6)$$

where $S' = \pm S$, the total angular momentum operators can be written as

$$\begin{aligned} J^2 &= -\frac{1}{\sin \theta} \partial_\theta (\sin \theta \partial_\theta) - \frac{1}{\sin^2 \theta} \partial_\phi^2 - \frac{i}{\sin^2 \theta} [(2S-1) \cos \theta - 2S'] \partial_\phi \\ &\quad + \frac{1}{4\sin^2 \theta} [(2S-1) \cos \theta - 2S']^2 + (S - \frac{1}{2})^2 \\ L^2 &= -\frac{1}{\sin \theta} \partial_\theta (\sin \theta \partial_\theta) - \frac{1}{\sin^2 \theta} \partial_\phi^2 - \frac{i}{\sin^2 \theta} [(2S+1) \cos \theta - 2S'] \partial_\phi \\ &\quad + \frac{1}{4\sin^2 \theta} [(2S+1) \cos \theta - 2S']^2 + (S + \frac{1}{2})^2 \end{aligned} \quad (7)$$

In the R_a region, we have the following relations ($S' = S$)

$$\begin{aligned} A\bar{A} &= J^2 + \frac{1}{4} - S^2 \\ \bar{A}A &= L^2 + \frac{1}{4} - S^2 \end{aligned} \quad (8)$$

In the R_b region, we have the same relations as Eq.(8) if one replaces $S' = -S$. The wave functions in these two regions have the following relations

$$\psi_{ai}(\theta, \phi) = e^{2iS\phi} \psi_{bi}(\theta, \phi), \quad i = 1, 2. \quad (9)$$

From Eq.(8), we can easily solve the eigenequation (4). In R_a region, we have the eigenfunctions

$$\begin{aligned} \psi_1(\theta, \phi) &= e^{i(m+S)\phi} Y_1^{Sjm}(\theta), \quad j = (S - \frac{1}{2}), (S - \frac{1}{2}) + 1, \dots, \quad m = -j, -j + 1, \dots, j \\ \psi_2(\theta, \phi) &= e^{i(n+S)\phi} Y_2^{Sln}(\theta), \quad l = (S + \frac{1}{2}), (S + \frac{1}{2}) + 1, \dots, \quad n = -l, -l + 1, \dots, l \end{aligned} \quad (10)$$

where $Y_1^{Sjm}(\theta)$ and $Y_2^{Sln}(\theta)$ are called the monopole harmonics [27], which satisfy the following equations

$$\begin{aligned} [j(j+1) - (S - \frac{1}{2})^2] Y_1^{Sjm}(\theta) &= \\ [-\frac{1}{\sin \theta} \partial_\theta (\sin \theta \partial_\theta) + \frac{1}{\sin^2 \theta} (m + (S - \frac{1}{2}) \cos \theta)^2] Y_1^{Sjm}(\theta) \\ [l(l+1) - (S + \frac{1}{2})^2] Y_2^{Sln}(\theta) &= \\ [-\frac{1}{\sin \theta} \partial_\theta (\sin \theta \partial_\theta) + \frac{1}{\sin^2 \theta} (n + (S + \frac{1}{2}) \cos \theta)^2] Y_2^{Sln}(\theta) \end{aligned} \quad (11)$$

The eigenvalues of Eq.(4) can be written as

$$\begin{aligned}
E_1^2 R^2 &= j(j+1) - (S - \frac{1}{2})(S + \frac{1}{2}), \quad j = (S - \frac{1}{2}) + n, \quad n = 0, 1, 2, \dots \\
E_2^2 R^2 &= l(l+1) - (S - \frac{1}{2})(S + \frac{1}{2}), \quad l = (S + \frac{1}{2}) + n, \quad n = 0, 1, 2, \dots
\end{aligned} \tag{12}$$

where the quantum numbers j and l satisfy the relation, $l = j + 1$. For $j = S - \frac{1}{2}$, we only have one set of wave functions $\psi_1(\theta, \phi) = e^{i(m+S)\phi} Y_1^{S, S-\frac{1}{2}, m}(\theta)$, $m = -(S - \frac{1}{2}), -(S - \frac{1}{2}) + 1, \dots, S - \frac{1}{2}$, corresponding to the eigenvalue $E_1(j = S - \frac{1}{2}) = 0$, called "zero-point" energy. For $j = S - \frac{1}{2} + n$, n is a non-zero integer, the eigenvalue $E_1(j) = E_2(l)$ is double degenerate for the wavefunctions $\psi_1(\theta, \phi)$ and $\psi_2(\theta, \phi)$. The lowest nonzero eigenvalue of the 2D Dirac fermion is ($j = S + \frac{1}{2}$)

$$E = (1 + \frac{1}{2S})^{1/2} \cdot \sqrt{2} E_B \tag{13}$$

where $E_B = \frac{e^2}{l_B}$ is the Coulomb-like energy of the 2D Dirac fermions, l_B is the magnetic length. For a free 2D Dirac fermion system, S is proportional to the number of the 2D Dirac fermions, according to the definition of the filling factor of the Landau levels, $\nu = \frac{\rho \Phi_0}{B}$, we can easily obtain the relation $2S = \frac{N}{\nu}$, so we have the total energy of N 2D Dirac fermions from Eq.(13)

$$E_N = \sqrt{2} N E_B + \frac{\sqrt{2}}{2} \nu E_B \tag{14}$$

We see that the last term is dependent upon the filling factor ν , which can be called as the "collective excitation energy" of the 2D Dirac fermion system, $\Delta E_N = \frac{\sqrt{2}}{2} \nu E_B$. For the filling factors $\nu = \frac{p}{2mp \pm 1}$, p and m are nonzero integers, the collective excitation energy ΔE_N takes discrete values for different p and m , and go to the limit $\Delta E_N(m, p \rightarrow \infty) = \frac{\sqrt{2}}{2} \cdot \frac{1}{2m} E_B$, for $p \rightarrow \infty$, which means that there exist infinite nearly degenerate levels around these energies for different m . It is reasonable to assume that the system has a continuous spectrum at these energies and show the Fermi-liquid-like behavior [9]. If we choose these energies as the original point of energy, we can obtain a collective excitation energy gap of the system for finite p

$$\Delta_\nu(m, p) = \mp \frac{\sqrt{2}}{2} \frac{E_B}{2m(2mp \pm 1)} \tag{15}$$

where the positive and negative signs correspond to the hole and fermion filling of the Landau levels. For a real material we should take $E_B = \frac{e^2}{\epsilon l_B}$, ϵ is the dielectric constant of the material. We see that the collective excitation energy gap is proportional to $\frac{1}{p}$, but proportional to $\frac{1}{m^2}$ for large m , this behavior is similarly consistent with the experimental data [12].

We compare the our exact results Δ_ν with the computing results in Ref. [32]. From Eq.(15), we have the following collective excitation energy gaps for different filling factors,

$$\begin{aligned}\Delta_{\frac{1}{3}}^{exc.} &\simeq 0.118E_B, & \Delta_{\frac{2}{5}}^{exc.} &\simeq 0.071E_B \\ \Delta_{\frac{3}{7}}^{exc.} &\simeq 0.051E_B, & \Delta_{\frac{2}{7}}^{exc.} &\simeq 0.025E_B\end{aligned}\tag{16}$$

while the computing results of Ref. [32] are that, $\Delta_{\frac{1}{3}}^c \simeq 0.102(3)E_B$, $\Delta_{\frac{2}{5}}^c \simeq 0.063(4)E_B$, $\Delta_{\frac{3}{7}}^c \simeq 0.049(18)E_B$, $\Delta_{\frac{2}{7}}^c \simeq 0.022(3)E_B$, they are very well agreement with our exact results. The small difference between them may derive from the computing results extrapoted from finite (a few electrons) system. To compare with the experimental data, we must redefine Eq.(15). For a fixed N (electron number) system, Eq.(15) can be changed into as when magnetic field B varies

$$\left(\frac{E_N}{E_B}\right)_{\nu=\frac{p}{2mp\pm 1}} - \left(\frac{E_N}{E_B}\right)_{\nu=\frac{1}{2m}} = \mp \frac{\sqrt{2}}{2} \cdot \frac{1}{2m(2mp \pm 1)}\tag{17}$$

This relation is exact for a non-disorder material. However, for an integer quantum Hall effect, if the composite fermion conception is valid, from Eq.(14) we see that the filling factor ν takes even or odd integer, the system may show different behavior; for $\nu = 2n$, taking even integer, we have the relation, $E_N(\nu = 2n) = E_{N+n}(\nu = 0)$, which means that we may have a zero collective excitation energy gap; while for $\nu = 2n + 1$, taking odd integer, we have the relation, $E_N(\nu = 2n + 1) = E_{N+n}(\nu = 1)$, which means we always have a nonzero collective excitation energy gap. If this picture of the integer quantum Hall effect is correct, the transitions[[33], [34]] between insulating state and the quantum Hall states for the filling factors $\nu = 1$ and $\nu = 2$ may be different, because for the former the collective excitation energy gap will influence this transition.

In summary, we have exactly solved the eigenequation of a 2D Dirac fermion moving on the surface of a sphere under the influence of a radial magnetic field B , and obtained an exact expression of the collective excitation energy gap for the filling factors $\nu = \frac{p}{2mp \pm 1}$, m and p are non-zero integers, which is very well agreement with the computing results.

We are very grateful to Prof. L. Yu and Prof. Z. B. Su for their encouragement.

*Present address: Max-Planck-Institut für Physik Komplexer Systeme, Bayreuther Str. 40, D-01187 Dresden, Germany.

REFERENCES

- [1] D.C.Tsui, H.Stormer, and A.Gossard, Phys. Rev. Lett. **48**, 1559(1982).
- [2] R.B Laughlin, Phys. Rev. Lett. **50**, 1395(1983).
- [3] J.K.Jain, Phys. Rev. Lett. **63**, 199(1989); Phys. Rev. **B40**, 8079(1989); **B41**, 7653(1990).
- [4] F.Wilczek, Phys. Rev. Lett. **48**, 1144(1982); **49**, 957(1982).
- [5] S.M.Girvin, and A.H.MacDonald, Phys. Rev. Lett. **58**, 1252(1987).
- [6] R.B Laughlin, Phys. Rev. Lett. **60**, 2677(1988); E.Rezayi, and F.D.M.Haldane, Phys. Rev. Lett. **61**, 1985(1988).
- [7] S.C.Zhang, H.Hanson, and S.Kivelson, Phys. Rev. Lett. **62**, 82(1989); N.Read, *ibid.*, **62**, 86(1989).
- [8] A.Lopez, and E.Fradkin, Phys. Rev. **B44**, 5246(1991); M.Greiter, X.G.Wen, and F.Wilczek, Phys. Rev. Lett. **66**, 3205(1991); G.Moore, and N.Read, Nucl. Phys. **B360**, 362(1991); S.Kivelson, D.H.Lee, and S.C.Zhang, Phys. Rev. **B46**, 2223(1992); D.Schmeltzer, *ibid.*, **B46**, 1591(1992).
- [9] B.I.Halperin, P.A.Lee, and N.Read, Phys. Rev. **B47**, 7312(1993); V.Kalmeyer, and S.C.Zhang, *ibid.*, **B46**, 9889(1992).
- [10] H.W.Jiang, *et al.*, Phys. Rev. **B40**, 12013(1989).
- [11] R.L.Willett, *et al.*, Phys. Rev. Lett. **65**, 112(1990); Phys. Rev. **B47**, 7344(1993).
- [12] R.R.Du, *et al.*, Phys. Rev. Lett. **70**, 2944(1993).
- [13] R.L.Willett, *et al.*, Phys. Rev. Lett. **71**, 3846(1993).
- [14] W.Kang, *et al.*, Phys. Rev. Lett. **71**, 3850(1993).
- [15] D.R.Leadly, *et al.*, Phys. Rev. Lett. **72**, 1906(1994).

- [16] V.J.Goldman, B.Su, and J.K.Jain, Phys. Rev. Lett. **72**, 2065(1994).
- [17] H.C.Manoharan, M.Shayegan, and S.J.Lepper, Phys. Rev. Lett. **73**, 3270(1994).
- [18] R.R.Du, *et al.*, Phys. Rev. Lett. **73**, 7274(1994).
- [19] B.Tieke, *et al.*, Phys. Rev. Lett. **76**, 3630(1996).
- [20] E.H.Rezayi, and A.H.MacDonald, Phys. Rev. **B44**, 8395(1991).
- [21] G.Dev, and J.K.Jain, Phys. Rev. Lett. **69**, 2843(1992).
- [22] X.G.Wu, G.Dev, and J.K.Jain, Phys. Rev. Lett. **71**, 153(1993).
- [23] E.Rezayi, and N.Read, Phys. Rev. Lett. **72**, 900(1994).
- [24] X.G.Wu, and J.K.Jain, Phys. Rev. **B51**, 1752(1995).
- [25] P.Sito, S.N.Yi, K.S.Yi, and J.J.Quinn, Phys. Rev. Lett. **76**, 3396(1996).
- [26] F.D.M.Haldane, Phys. Rev. Lett. **51**, 605(1983).
- [27] T.T.Wu, and C.N.Yang, Nucl. Phys. **B107**, 365(1976); Phys. Rev. **D16**, 1018(1977).
- [28] G.Fano, F.Ortolani, and E.Colombo, Phys. Rev. **B34**, 2670(1986).
- [29] Y.L.Liu, Phys. Lett. **A212**, 103(1996).
- [30] S.Deser, R.Jackiw, and S.Templeton, Phys. Rev. Lett. **48**, 975(1982); Ann. Phys. (N.Y.) **140**, 372(1982).
- [31] N.D.Birrel, and P.C.W.Davies, *Quantum Fields in Curved Space*, (Cambridge Univ. Press, London, 1982).
- [32] N.d'Ambrumenil, and R.Morf, Phys. Rev. **B40**, 6108(1989).
- [33] H.W.Jiang, *et al.*, Phys. Rev. Lett. **65**, 633(1990); **71**, 1439(1993); R.Hughes, *et al.*, J. Phys.: Condens. Matter **6**, 4763(1994).

[34] I.Glozman, *et al.*, Phys. Rev. Lett. **74**, 594(1995); D.Shahar, *et al.*, *ibid.*, **74**, 4511(1995).